

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 January 2004 (22.01.2004)

PCT

(10) International Publication Number
WO 2004/007454 A1

(51) International Patent Classification⁷: C07D 211/46,
405/06, 401/06, 413/06, A61K 31/445, A61P 3/00, 35/00,
9/00

(74) Agent: BLAKEY, Alison, Jane; Oxford GlycoSciences
(UK) Ltd, The Forum, 86 Milton Park, Abingdon, Oxford-
shire OX14 4RY (GB).

(21) International Application Number:
PCT/GB2003/003244

(22) International Filing Date: 17 July 2003 (17.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0216656.9 17 July 2002 (17.07.2002) GB
0301480.0 22 January 2003 (22.01.2003) GB
0313674.4 13 June 2003 (13.06.2003) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): OXFORD
GLYCOSCIENCES (UK) LTD [GB/GB]; The Forum, 86
Milton Park, Abingdon, Oxfordshire OX14 4RY (GB).


Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): ALI, Mezher, Hus-
sein [GB/GB]; The Forum, 86 Milton Park, Abingdon, Ox-
fordshire OX14 4RY (GB). ORCHARD, Michael, Glen
[GB/GB]; The Forum, 86 Milton Park, Abingdon, Oxford-
shire OX14 4RY (GB).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 2004/007454 A1

(54) Title: PIPERIDINETRIOL DERIVATIVES AS INHIBITORS OF GLYCOSYLCERAMIDS SYNTHASE



(I)

(57) Abstract: Compounds of formula (I); wherein R represents various substituent groups, are useful as inhibitors of glucosylce-
ramide synthase.

PIPERIDINETRIOL DERIVATIVES AS INHIBITORS OF GLYCOSYLCERAMIDSYNTHASE

The present invention relates to novel piperidine derivatives useful as inhibitors of glucosylceramide synthase (GCS; UDP-glucose:ceramide glucosyltransferase, UDP-glucose:N-acylsphingosine D-glucosyltransferase, EC 2.4.1.80), methods for their preparation and their use in medicine, specifically in the treatment and prevention of disease states mediated by GCS. The compounds find use in the treatment of glycolipid storage diseases, diseases associated with glycolipid accumulation, cancers in which glycolipid synthesis is abnormal, infectious diseases caused by organisms which use cell surface glycolipids as receptors, infectious diseases in which synthesis of glucosylceramide is essential or important, diseases in which excessive glycolipid synthesis occurs, neuronal disorders, neuronal injury and inflammatory diseases or disorders associated with macrophage recruitment and activation.

GCS is an intracellular enzyme that catalyzes the assembly of uridine diphosphate-glucose and ceramide into the glycolipid, glucosylceramide. The role of GCS in regulating ceramide levels has been explored, since this molecule can induce apoptotic cell death (J. Biol. Chem., 2000, 275(10), 7138-43). The role of GCS in maintaining cholesterol/glycolipid 'rafts', cell-surface membrane domains of specialized permeability and functionality that appear to be involved in a variety of signal transduction events, has also been investigated (Nature, 1997, 387(6633), 569-72).

GCS is considered to be a target for treating certain human diseases. Glucosylceramide and structurally related glycolipids are stored in the lysosomes of patients with genetic diseases, which result from a mutation in one of the essential glycolipid-degrading enzymes (e.g. Gaucher, Tay Sachs, Sandhoffs, GM1 gangliosidosis and Fabry diseases). Glycolipid storage also occurs as a secondary effect in some tissues (e.g. neuronal tissue) with genetic storage diseases such as Niemann-Pick C disease, mucopolysaccharidoses, mucopolipidosis type IV (Proc. Natl. Acad. Sci. USA, 1998, May 26, 95(11), 6373-8) and α -mannosidosis (Proc. Natl. Acad. Sci. USA, 1991, Dec 15, 88(24), 11330-4). GCS inhibitors may be applied to reduce the rate of glycolipid synthesis in diseased cells so that there is less glycolipid present to be stored, a treatment approach termed substrate deprivation. Studies have demonstrated that GCS inhibitors can be used to reduce the glycolipid accumulation seen in cell and animal models of glycolipid storage disorders (Proc. Natl. Acad. Sci. USA, 1999, 96(11), 6388-93; Science, 1997, 276(5311), 428-31; J. Clin. Invest., 2000, 105(11), 1563-71). Furthermore, clinical trials have shown that GCS inhibitors, such as, N-butyldeoxynojirimycin (NB-DNJ) are useful in treating human patients with Gaucher disease (Lancet, 2000, 355(9214), 1481-5). The use of the imino sugar NB-DNJ as a GCS inhibitor is disclosed in EP-A-0698012. EP-A-0536402 and EP-A-0698012 disclose that N-alkyl derivatives of deoxygalactonojirimycin, e.g. N-butyldeoxygalactonojirimycin (NB-DGJ), may also be of use in the treatment of glycolipid storage disorders. EP-A-0698012 also discloses that the corresponding N-butyl derivatives of mannose (NB-DMJ), fucose (NB-DFJ) and N-acetylglucosamine (NB-NAG) do not act as inhibitors of glycolipid biosynthesis.

The use of GCS inhibitors in the treatment of human malignancies has also been proposed. Tumours can synthesize abnormal quantities of glycolipids that are typically present/ absent in normal tissues. In addition glycolipids, or gangliosides, in particular are shed by tumour cells and released into the extracellular space and the bloodstream. Both tumour shed and cell surface bound tumour gangliosides can influence tumour host cell interactions such as cell-cell contacts or adhesion (Methods Enzymol., 2000, 312, 447-58), cell motility (Mol. Chem. Neuropathol., 1995, 24(2-3), 121-35), growth factor signalling events (J. Biol. Chem., 2000, 275(44), 34213-23), tumour stimulated angiogenesis (Acta. Oncol., 1997, 36(4), 383-7) and tumour specific immune responses (J. Immunol., 1999, Oct 1, 163(7), 3718-26). All these events can affect tumour development and progression. Glycolipids, glucosylceramide in particular, are known to accumulate in multidrug resistant (MDR) tumour cells

(Anticancer Res., 1998, 18(1B), 475-80) and in vitro treatment of these cells with GCS inhibitors can reverse the MDR phenotype (J. Biol. Chem., 1997, 272(3), 1682-7; Br. J. Cancer, 1999, 81(3), 423-30).

Cell surface glycolipids also have roles in infectious disease, serving as receptors for the binding of pathogenic bacteria (APMIS, 1990, Dec, 98(12), 1053-60, Review), fungi (Infect. Immun., 1990 Jul, 58(7), 2085-90) and viruses (FEBS Lett., 1984, May 7, 170(1), 15-18). In addition, glycolipids on the surface of cells are bound by bacterial toxins (Methods Enzymol., 2000, 312, 459-73) for instance, the B subunit of cholera toxin (ganglioside GM1) and verocytotoxin (globotriaosylceramide GB3) (J. Infect. Dis., 2001, suppl. 70-73, 183).

GCS inhibitors may also find use in the treatment of viral infections.

The use of GCS inhibitors may also be appropriate in a number of other clinical indications which are associated with abnormalities in glycolipid synthesis. Atherosclerotic lesions of human aorta have a higher ganglioside content than unaffected regions of the aorta and serum ganglioside concentrations in atherosclerotic patients are higher than in normal individuals (Lipids, 1994, 29(1), 1-5). Tissue derived from the kidneys of patients with polycystic kidney disease contains high levels of both glucosylceramide and lactosylceramide (J. Lipid. Res., 1996, Jun, 37(6), 1334-44). Renal hypertrophy in an animal model of diabetes is associated with increases in glycolipid synthesis, (J. Clin. Invest., 1993, Mar, 91(3), 797-803).

Glycolipid metabolism also plays a critical role in neuronal disorders, such as Alzheimer's disease and epilepsy. For instance, Niemann-Pick C (NPC) patient neurons present with fibrillar tangles reminiscent of the morphology seen in Alzheimer's disease.

GM1 ganglioside binding by amyloid beta-protein induces conformational changes that support its formation of fibrous polymers and the fibrillar deposition of this protein is an early event in Alzheimer's disease (Yanagisawa et al., 1995, Nat. Med. 1, 1062-6; Choo-Smith et al., 1997, Biol. Chem., 272, 22987-90). Thus, decreasing GM1 synthesis by using agents such as GCS inhibitors, e.g. NB-DNJ, could inhibit the fibre formation seen in Alzheimer's disease.

In contrast, preliminary clinical trials have shown that neurodegenerative processes seen in Parkinson's disease, stroke and spinal cord injuries seem to improve by treating patients with GM1 ganglioside (Alter, (1998), Ann. NY Acad. Sci., 845, 391-4011; Schneider, 1998, Ann. NY. Acad. Sci., 845, 363-73; Geisler, (1998), Ann. NY. Acad. Sci., 845, 374-81). It is possible that co-administering glucosylceramide synthesis inhibitors would provide the clinician greater control over this treatment course. GCS inhibitors like NB-DNJ would limit patient-specific inconsistencies by blocking their neuronal glycolipid synthesis. In addition, inhibiting glucosylceramide synthesis would limit the metabolism of administered glycolipids into other, perhaps unproductive, forms. Thus, the ability to modulate glucosylceramide synthesis with GCS inhibitors may be useful in the treatment of a wide variety of neuronal disorders.

In addition, it has also been shown that imino sugars can reversibly induce male sterility and can, therefore, be used as male contraceptives. Also, GCS inhibitors could be used for the treatment of obesity.

A role for glycolipids in some aspects of inflammatory or immune responses has also been suggested. Following an inflammatory stimulus, such as that obtained with thioglycolate, the ganglioside profile of murine peritoneal macrophages changes from a simple profile (3 major species) in resting macrophage to a more complex profile (more than 14 species) in activated and recruited macrophage, see Ryan, J.L. et al., Yale J. Biol. Med., 1985, 58(2) 125-31; Yohe, H.C. et al., Biochim. Biophys. Acta., 1985, 818(1), 81-6; Yohe, H.C. et al., Immunol., 1991, 146(6), 1900-8. Furthermore, *in vivo* administration of an inflammatory agent, e.g. bacterial endotoxin, results in the increased expression of two enzymes, serine palmitoyltransferase and glucosylceramide synthase, which are key to the *de novo* synthesis of glycolipids, see Memon, R.A. et al., J. Biol. Chem., 1999, 274(28), 19707-13; Memon, R.A.

et al., J. Lipid. Res., 2001, 42(3), 452-9.

Such a role for glycolipids is further supported by the demonstration of changes in glycolipid expression in animals with genetic defects which result in hyper- or hypo- sensitive responses to inflammatory stimuli. For example, upon endotoxin treatment in C3H/HeJ mice, which have a toll-like receptor 4 mutation and are hypo-responsive to bacterial endotoxin, recruited macrophages were found to lack ganglioside G_{M1b} , which is a major ganglioside found in recruited macrophages in normal mice, see Yohe, H.C. et al., Immunol., 1991, 146(6), 1900-8; Yohe, H.C. et al., Immunol., 1986, 137(12), 3921-7.


Hence, GCS inhibitors may be useful in the treatment of inflammatory diseases and other disorders associated with macrophage recruitment and activation, including but not limited to, rheumatoid arthritis, Crohn's disease, asthma and sepsis.

WO02/055498, published after the priority date of the present application, discloses piperidine derivatives useful as GCS inhibitors.

Given the importance of GCS in a wide spectrum of diseases, it is essential that new tools that provide a means for modulating this enzyme's function be developed. Towards this end, we have identified a class of novel compounds that are useful in inhibiting GCS's catalytic activity.

The compounds of the invention may exhibit improved potency and/or selectivity for GCS, relative to non-lysosomal- β -glucocerebrosidase activity, over known hydroxylated piperidine derivatives.

According to the invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt or prodrug thereof:



wherein

R is C_{1-3} alkylAr¹ where Ar¹ is phenyl or pyridyl;

wherein phenyl is substituted by one or more substituents selected from CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R⁵)₂, N(R¹)COR², N(R¹)SO₂R², C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one or more of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O, S and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C₁₋₆ alkyl and C₀₋₃ alkylAr⁴;

and the Ar¹ phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF₃, OCF₃, OR³ and C₁₋₆ alkyl;

and wherein pyridyl is substituted by one or more substituents, selected from, CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R⁵)₂, N(R¹)COR², N(R¹)SO₂R², F, Cl, Br, CF₃, OCF₃, OR³, C₁₋₆ alkyl, C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O, S and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C₁₋₆ alkyl and C₀₋₃ alkylAr⁴;

R^1 is H, C_{1-6} alkyl optionally substituted by OH, Ar^3 , or C_{1-6} alkyl Ar^3 , or the group $N(R^1)_2$ may form a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O, S and NR^3 and is optionally substituted by an oxo group;

R^2 is C_{1-6} alkyl optionally substituted by OH, Ar^3 , or C_{1-6} alkyl Ar^3 ;

5 R^3 is H, or C_{1-6} alkyl;

R^4 is H, C_{1-6} alkyl or C_{0-3} alkyl Ar^4 ;

R^5 is H, C_{1-6} alkyl optionally substituted by OH, Ar^3 or C_{1-6} alkyl Ar^3 , or the group $N(R^5)_2$ may form a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O, S and NR^3 and is optionally substituted by an oxo group;

10 Ar^2 and Ar^3 are independently phenyl or a 5- to 10-membered heteroaryl group containing up to 3 heteroatoms selected from O, S and NR^3 , which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl;

Ar^4 is phenyl or pyridyl either of which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl; and

15 n is 0, 1 or 2.

R is preferably C_1 alkyl Ar^1 .

Ar^1 is preferably phenyl, wherein phenyl is substituted as defined for formula (I).

Ar^1 phenyl is preferably substituted on the para position.

20 More preferably Ar^1 is phenyl, wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, $SO_2N(R^1)_2$, $N(R^5)_2$, $N(R^1)COR^2$, C_{0-6} alkyl Ar^2 and C_{2-6} alkenyl Ar^2 wherein one or more of the $-CH_2-$ groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR^3 , provided that when the heteroatom is O, at least two $-CH_2-$ groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar^1 phenyl may together form a fused 5- or 6-

25 membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O and NR^4 and is optionally substituted by one or more substituents selected from, an oxo group, C_{1-6} alkyl and C_{0-3} alkyl Ar^4 , and the Ar^1 is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

Yet more preferably Ar^1 is phenyl, wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, $N(R^5)_2$ and C_{0-6} alkyl Ar^2 wherein one or more of the $-CH_2-$ groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR^3 , provided that when the heteroatom is O, at least two $-CH_2-$ groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar^1 phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O and NR^4 and is

35 optionally substituted by one or more substituents selected from, an oxo group, C_{1-6} alkyl and C_{0-3} alkyl Ar^4 , and the Ar^1 is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

Even more preferably Ar^1 is phenyl, wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, $N(R^5)_2$ and C_{0-6} alkyl Ar^2 wherein one or more of the $-CH_2-$ groups of the alkyl chain may be replaced with O, provided that at least two $-CH_2-$ groups separate it from any additional O atom introduced into the alkyl chain, and the Ar^1 phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

40

When Ar^1 is phenyl and has an additional optional substituent as defined for formula (I) on the ortho position, the substituent is preferably selected from OCH_3 and F. More preferably the ortho substituent is F.

45

When Ar^1 is phenyl substituted by C_2 alkyl Ar^2 wherein one of the $-CH_2-$ groups of the alkyl chain is replaced with O, preferably, the $-CH_2-$ group linked to the Ar^1 phenyl is replaced with O.

R^1 is preferably H, C_{1-6} alkyl or C_{1-6} alkyl Ar^3 . More preferably R^1 is H or C_{1-6} alkyl Ar^3 .

R^2 is preferably Ar^3 or C_{1-6} alkyl Ar^3 . More preferably R^2 is C_{1-6} alkyl Ar^3 .

5 R^3 is preferably H.

R^4 is preferably H or C_{1-6} alkyl. More preferably R^4 is H.

R^5 is preferably C_{1-6} alkyl optionally substituted by OH, or C_{1-6} alkyl Ar^3 . More preferably R^5 is C_{1-6} alkyl.

10 For the groups R^1 , R^2 or R^5 , the group C_{1-6} alkyl Ar^3 is preferably C_{1-3} alkyl Ar^3 , for example, C_1 alkyl Ar^3 or C_2 alkyl Ar^3 .

Ar^2 is preferably phenyl which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

Ar^3 is preferably phenyl which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

15 Ar^4 is preferably phenyl which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl.

n is preferably 2.

In the groups $CON(R^1)_2$, $SO_2N(R^1)_2$, and $N(R^5)_2$ the R^1 and R^5 groups may be the same or different.

20 When two adjacent substituents on the Ar^1 form a fused 5- or 6-membered saturated or unsaturated ring optionally containing 1 or 2 heteroatoms selected from O, S and NR⁴, examples of bicyclic groups which may be formed include benzofuran, indole, benzoxazine, quinoline and isoquinoline.

When $N(R^1)_2$ forms a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O, S and NR³, examples of heterocyclic groups include, piperidine, piperazine, morpholine and quinoline.

When $N(R^5)_2$ forms a 5- to 10-membered heterocyclic group, preferably a 5- or 6-membered heterocyclic group, optionally containing one or more additional heteroatoms selected from O, S and NR³, examples of heterocyclic groups include, piperidine, piperazine and morpholine.

30 When Ar^2 or Ar^3 is a 5- to 10-membered heteroaryl group, examples of heteroaryl groups include furan, thiophene, oxazole, triazole, pyridine, pyrazine, pyrimidine, benzofuran benzothiophene and benzoxazine.

A specific group of compounds of formula (I) which may be mentioned are those where R is C_{1-3} alkylaryl where aryl is phenyl or pyridyl;

35 wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, SO_2R^2 , $N(R^2)_2$, and $N(R^1)COR^2$, or two adjacent substituents on the phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring optionally containing 1 or 2 heteroatoms, selected from N and O, and the phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , and OR^1 ;

40 and wherein pyridyl is substituted by one or more substituents, selected from, CN, $CON(R^1)_2$, SO_2R^2 , $N(R^2)_2$, $N(R^1)COR^2$, F, Cl, Br, CF_3 , OCF_3 , and OR^1 or two adjacent substituents on the pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring which may optionally contain 1 or 2 heteroatoms, selected from N and O;

45 R^1 is H, or C_{1-6} alkyl; and

R^2 is C_{1-6} alkyl.

Another specific group of compounds of formula (I) that may be mentioned are those where R is C_{1-3} alkylAr¹ where Ar¹ is phenyl or pyridyl;

wherein phenyl is substituted by one or more substituents selected from CN, CON(R¹)₂, SO₂R², N(R²)₂, N(R¹)COR², and C_{1-3} alkylAr² wherein one of the -CH₂- groups of the alkyl chain may be replaced with O and where Ar² is phenyl or a 5- or 6-membered heteroaryl group containing up to 3 heteroatoms selected from N, O and S; or two adjacent substituents on the Ar¹ phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring optionally containing 1 or 2 heteroatoms selected from N and O;

and wherein pyridyl is substituted by one or more substituents, selected from, CN, CON(R¹)₂, SO₂R², N(R²)₂, N(R¹)COR², F, Cl, Br, CF₃, OCF₃, OR³, and C_{1-3} alkylAr² wherein one of the -CH₂- groups of the alkyl chain may be replaced with O and where Ar² is phenyl or a 5- or 6-membered heteroaryl group containing up to 3 heteroatoms selected from N, O and S; or two adjacent substituents on the pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring optionally containing 1 or 2 heteroatoms selected from N and O;

and the Ar¹ or Ar² phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF₃, OCF₃, and OR³;

R¹ is H, C_{1-6} alkyl, Ar³, or C_{1-6} alkylAr³ where Ar³ is phenyl or a 5- or 6-membered heteroaryl group containing up to 3 heteroatoms selected from N, O and S, any of which may be substituted by one or more substituents selected from F, Cl, Br, CF₃, OCF₃, and OR³;

R² is C_{1-6} alkyl, Ar³, or C_{1-6} alkylAr³ where Ar³ is phenyl or a 5- or 6-membered heteroaryl group containing up to 3 heteroatoms selected from N, O and S, any of which may be substituted by one or more substituents selected from F, Cl, Br, CF₃, OCF₃, and OR³; and

R³ is H, or C_{1-6} alkyl.

Yet another specific group of compounds of formula (I) that may be mentioned are those where R is C_{1-3} alkylAr¹ where Ar¹ is phenyl or pyridyl;

wherein phenyl is substituted by one or more substituents selected from CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R²)₂, N(R¹)COR², N(R¹)SO₂R², C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one or more of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C_{1-6} alkyl and C₀₋₃ alkylAr⁴;

and the Ar¹ phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF₃, OCF₃, OR³ and C_{1-6} alkyl;

and wherein pyridyl is substituted by one or more substituents, selected from, CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R²)₂, N(R¹)COR², N(R¹)SO₂R², F, Cl, Br, CF₃, OCF₃, OR³, C_{1-6} alkyl, C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C_{1-6} alkyl and C₀₋₃ alkylAr⁴;

R¹ is H, C_{1-6} alkyl, Ar³, or C_{1-6} alkylAr³, or the group N(R¹)₂ may form a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O and NR³;

R^2 is C_{1-6} alkyl, Ar^3 , or C_{1-6} alkyl Ar^3 , or the group $N(R^2)_2$ may form a 5- to 6-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O and NR^3 and is optionally substituted by an oxo group;

R^3 is H, or C_{1-6} alkyl;

5 R^4 is H, C_{1-6} alkyl or C_{0-3} alkyl Ar^4 ;

Ar^2 is phenyl or a 5- or 6-membered heteroaryl group containing up to 3 heteroatoms selected from O, S and NR^3 , any of which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl;

10 Ar^3 is phenyl or a 5- to 10-membered heteroaryl group containing up to 3 heteroatoms selected from O, S and NR^3 , any of which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl;

Ar^4 is phenyl or pyridyl either of which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and C_{1-6} alkyl; and

n is 0, 1 or 2.

15

The compounds of the invention preferably have a molecular weight of less than 800, more preferably less than 600.

20 The term "alkyl" as used herein whether on its own or as part of a larger group e.g. "alkylaryl", includes both straight and branched chain radicals. The term alkyl also includes those radicals wherein one or more hydrogen atoms are replaced by fluorine. Alkenyl and alkynyl are to be interpreted accordingly.

The term "heterocyclic group" as used herein includes, unless otherwise defined, non-aromatic and aromatic, single and fused, rings containing one or more, e.g. up to three, heteroatoms in each ring, each of which is selected from O, S and N, which rings, may be unsubstituted or substituted. Each
25 heterocyclic ring suitably has from 5 to 10, preferably 5, 6, 9 or 10 ring atoms. A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring. Examples of heterocyclyl groups, including heteroaromatic ring systems, are as follows: pyrrolidine, piperidine, piperazine, morpholine, imidazolidine, pyrazolidine, pyrrole, quinoline, isoquinoline, pyridine, pyrazine, pyrimidine, oxazole, thiazole, thiophene, indole, furan, thiadiazole, triazole, imidazole, benzopyran,
30 benzofuran, benzothiophene, benzoxazine and benzamidazole. "Heteroaryl" is to be interpreted accordingly.

Specific compounds of the invention include the compounds provided in the Examples and pharmaceutically acceptable salts and prodrugs thereof.

Preferred compounds of the invention include:

35 Benzamide, N-[(4-fluorophenyl)methyl]-4-[[[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,

3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[[4-(phenylmethoxy)phenyl]methyl]-, (2S,3S,4R,5S),

Benzamide, N-[1-(R)-(phenyl)ethyl]-4-[[[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-;

40 3,4,5-Piperidinetriol, 1-[(3-cyano-4-(dipropylamino)phenyl)methyl]-2-(hydroxymethyl)-, (2S,3S,4R,5S),

Benzamide, N-[1-(S)-(4-fluorophenyl)ethyl]-4-[[[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,

45 Benzamide, N-[1-(R)-(4-fluorophenyl)ethyl]-4-[[[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-;

3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[(2-phenyl-2H-1,4-benzoxazin-3(4H)-one-6-yl)methyl]-, (2S,3S,4R,5S) and pharmaceutically acceptable salts and prodrugs thereof.

A highly preferred compound of the invention is:

3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[[4-(phenylmethoxy)phenyl]methyl]-, (2S,3S,4R,5S) and pharmaceutically acceptable salts and prodrugs thereof.

As described herein, for all aspects of the invention, reference to compounds of formula (I) encompasses pharmaceutically acceptable salts and prodrugs thereof.

As described herein, the compounds of the present invention can be used for the inhibition of GCS. Thus, an aspect of the present invention provides the use of the compounds of formula (I) in medicine.

Suitable, pharmaceutically acceptable salts of the compounds of formula (I) include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide and nitrate, or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, palmitate, salicylate and stearate.

Suitable prodrugs of the compounds of formula (I) include, but are not limited to, pharmaceutically acceptable esters such as C₁₋₆ alkyl esters.

Some of the compounds of this invention may be crystallised or recrystallised from solvents such as aqueous and organic solvents. In such cases solvates may be formed. This invention includes within its scope stoichiometric solvates including hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation.

Certain of the R groups of compounds of formula (I) may exist in the form of optical isomers, e.g. diastereoisomers and mixtures of isomers in all ratios, e.g. racemic mixtures. The invention includes all such forms, in particular the pure isomeric forms. The different isomeric forms may be separated or resolved one from the other by conventional methods, or any given isomer may be obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.


The compounds of the invention may exist as tautomers, e.g. keto/enol tautomers, all of which are included within the scope of formula (I).

Since the compounds of formula (I) are intended for use in pharmaceutical compositions it will readily be understood that they are preferably provided in substantially pure form, for example at least 60% pure, more suitably at least 75% pure and preferably at least 85%, especially at least 98% pure (% are on a weight for weight basis). Impure preparations of the compounds may be used for preparing the more pure forms used in the pharmaceutical compositions; these less pure preparations of the compounds should contain at least 1%, more suitably at least 5%, e.g. 10 to 59%, of a compound of formula (I) or pharmaceutically acceptable derivative thereof.


The compounds of formula (I) can be prepared by art-recognized procedures from known or commercially available starting materials. If the starting materials are unavailable from a commercial source, their synthesis is described herein, or they can be prepared by procedures known in the art.

Specifically, the compounds of formula (I) may be prepared by processes comprising:

a) reductive amination of an aldehyde of formula R⁵CHO wherein R⁵ is C₀₋₂ alkylAr¹ where Ar¹ is as defined in formula (I) with a compound of formula (II):




the reductive amination may be performed by methods known to those skilled in the art, e.g. using NaBH_3CN or a supported reagent such as (polystyrylmethyl) trimethylammonium cyanoborohydride in acetic acid-methanol or HCl -methanol, or using $\text{NaBH}(\text{OAc})_3$ in a solvent, such as dichloromethane; or
 b) deprotection of a compound of formula (III):



wherein R is as defined in formula (I) and P, which may be the same or different, are hydroxy protecting groups e.g. benzyl or substituted benzyl. When P is benzyl or substituted benzyl the deprotection is preferably conducted in the presence of hydrogen gas and a catalyst, such as, PdCl_2 or palladium on carbon in a suitable solvent, such as, an alcohol, e.g. ethanol. It will be understood that when P is benzyl or substituted benzyl and R is substituted benzyl, the R group can also be removed under these conditions to give compounds of formula (II). Thus, compounds of formula (I) where R is substituted benzyl are preferably produced using process a) above.


The compound of formula (II) is known, see e.g. Tet. Lett., 1997, 38(45), 8009-12.

Compounds of formula (III) may be prepared by reacting a compound of formula (IV):



wherein OL, which may be the same or different are leaving groups, such as mesyloxy, and P is as defined for formula (III), with an amine of formula RNH_2 , wherein R is as defined in formula (I), either neat or in a solvent such as tetrahydrofuran.

Compound (IVa), wherein L is mesyl and P is benzyl, may be prepared by reacting 2,3,4,6-tetra-O-benzyl-D-galactitol with mesyl chloride in the presence of a base such as pyridine, as described in WO02/055498.



Any novel intermediate compounds as described herein also fall within the scope of the present invention. Thus according to a further aspect of the invention there is provided a compound of formula (III) as defined above.

The invention also provides a compound of formula (I) when produced according to the methods described above.

During the synthesis of the compounds of formula (I) labile functional groups in the intermediate compounds, e.g. hydroxy, carboxy and amino groups, may be protected. A comprehensive discussion of the ways in which various labile functional groups may be protected and methods for cleaving the resulting protected derivatives is given in for example *Protective Groups in Organic Chemistry*, T.W. Greene and P.G.M. Wuts, (Wiley-Interscience, New York, 2nd edition, 1991).

The compounds of formula (I) may be prepared singly or as compound libraries comprising at least 2, for example 5 to 500 compounds and more preferably 10 to 100 compounds of formula (I).

Libraries of compounds of formula (I) may be prepared by multiple parallel synthesis using either solution phase or solid phase chemistry, by procedures known to those skilled in the art.

Thus according to a further aspect of the invention there is provided a compound library comprising at least 2 compounds of formula (I) or pharmaceutically acceptable salts or prodrugs thereof.

5 The pharmaceutically acceptable salts and prodrugs of the compounds of formula (I) may be prepared by methods well known to those skilled in the art.

The pharmaceutically effective compounds of formula (I) may be administered in conventional dosage forms prepared by combining a compound of formula (I) ("active ingredient") with standard pharmaceutical carriers, excipients or diluents according to conventional procedures well known in the art. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation.

According to a further aspect, the present invention provides a pharmaceutical composition comprising a compound of formula (I), together with one or more pharmaceutically acceptable carriers, excipients and/or diluents.

15 The active ingredient or pharmaceutical composition can be administered simultaneously, separately or sequentially with another treatment for the disorder to be treated.

The active ingredient or pharmaceutical composition may be administered to a subject by any of the routes conventionally used for drug administration, for example they may be adapted for oral (including buccal, sublingual), topical (including transdermal), nasal (including inhalation), rectal, vaginal or parenteral (including subcutaneous, intramuscular, intravenous or intradermal) administration to mammals including humans. The most suitable route for administration in any given case will depend on the particular compound or pharmaceutical composition, the subject and the nature and severity of the disease and the physical condition of the subject. Such compositions may be prepared by any method known in the art of pharmacy, for example by bringing into association the active ingredient with the carrier(s), excipient(s) and/ or diluent(s).

25 Pharmaceutical compositions adapted for oral administration may be presented as discrete units such as capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-aqueous liquids; edible foams or whips; or oil-in-water liquid emulsions or water-in-oil liquid emulsions.

Tablets and capsules for oral administration may be in unit dose presentation form and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants, for example potato starch; or acceptable wetting agents such as sodium lauryl sulfate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives, such as suspending agents, for example sorbitol, methyl cellulose, glucose syrup, gelatin, hydroxyethyl cellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and, if desired, conventional flavouring or colouring agents.

45 Pharmaceutical compositions adapted for topical administration may be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, impregnated dressings, sprays, aerosols or oils and may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration

and emollients in ointments and creams. Such applications include those to the eye or other external tissues, for example the mouth and skin and the compositions are preferably applied as a topical ointment or cream. When formulated in an ointment, the active ingredient may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredient may be formulated in a cream with an oil-in-water cream base or a water-in-oil base. The composition may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions.

Pharmaceutical compositions adapted for topical administration to the eye include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent.

Pharmaceutical compositions adapted for topical administration in the mouth include lozenges, pastilles and mouth washes.

Pharmaceutical compositions adapted for transdermal administration may be presented as discrete patches intended to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. For example, the active ingredient may be delivered from the patch by iontophoresis as generally described in Pharmaceutical Research, 3(6), 318, (1986).

Pharmaceutical compositions adapted for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns which is administered by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable compositions wherein the carrier is a liquid, for administration as a nasal spray or as nasal drops, include aqueous or oil solutions of the active ingredient.

Pharmaceutical compositions adapted for administration by inhalation include fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulizers or insufflators.

Pharmaceutical compositions adapted for rectal administration may be presented as suppositories or enemas. Suppositories will contain conventional suppository bases, e.g. cocoa-butter or other glyceride:

Pharmaceutical compositions adapted for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray compositions.

Pharmaceutical compositions adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets.

For parenteral administration, fluid unit dosage forms are prepared utilizing the active ingredient and a sterile vehicle, e.g. water. The active ingredient, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the active ingredient can be dissolved in water for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

Advantageously, agents such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. To enhance stability, the composition can be frozen after filling into the vial and the water removed under vacuum. The dry lyophilized powder is then sealed in the vial and an accompanying vial of water for injection may be supplied to reconstitute the liquid prior to use. Parenteral suspensions are prepared in substantially the same manner except that the active ingredient is suspended in the vehicle instead of being dissolved and sterilization cannot be accomplished by filtration. The active ingredient can be sterilised by exposure to ethylene oxide before suspending in the sterile vehicle.

Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the active ingredient.

The pharmaceutical compositions according to the invention are preferably adapted for oral administration.

5 It should be understood that in addition to the ingredients particularly mentioned above, the compositions may also include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents. They may also contain therapeutically active agents in addition to the compounds of the present invention. Such carriers may be present as from about 1% up to about 98% of the formulation. More usually they will form up to
10 about 80% of the formulation.

The compositions may contain from 0.1% by weight, e.g. 10-60% by weight, of the active material, depending on the method of administration.

Pharmaceutical compositions may be presented in unit dose forms containing a predetermined amount of active ingredient per dose. Such a unit may contain for example 0.1mg/kg to 750mg/kg, more
15 preferably 0.1mg/kg to 10mg/kg depending on the condition being treated, the route of administration and the age, weight and condition of the patient. Preferred unit dosage compositions are those containing a daily dose or sub-dose, or an appropriate fraction thereof, of an active ingredient.

It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of active ingredients will be determined by the nature and extent of the condition being treated, the
20 form, route and site of administration and the particular subject being treated and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e. the number of doses of the active ingredients given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

No toxicological effects are indicated when the compounds of formula (I) are administered in the
25 above mentioned dosage range.

The compounds of the invention are useful in that they are capable of inhibiting glucosylceramide synthase. Thus, the compounds of the invention can be used in the treatment of various glycolipid storage diseases, such as, Gaucher's disease, Sandhoffs disease, Tay-Sachs disease, Fabry disease, GM1
30 gangliosidosis etc. In addition, compounds, such as, this also can find use in the treatment of conditions in which glycolipid accumulation occurs, such as, Niemann-Pick disease, mucopolysaccharidoses (MPS I, MPS IIIA, MPS IIIB, MPS VI and MPS VII, preferably MPS I), mucopolidosis type IV and α -mannosidosis.

The compounds of the present invention can also be used in the treatment of cancers in which glycolipid synthesis is abnormal, such as, brain tumours, neuroblastoma, malignant melanoma, renal adenocarcinoma and multi-drug resistant cancers in general.

35 The compounds of the present invention can also be used in the treatment of diseases caused by infectious organisms which use cell surface glycolipids as receptors for either the infectious organism itself or for a toxin produced by the infectious organism (e.g. for attachment and/or invasion onto/into the host cell).

The compounds of the present invention may also be used in the treatment of diseases caused by infectious organisms for which the synthesis of glucosylceramide is an essential or important process, such
40 as, pathogenic fungi, e.g. *Cryptococcus neoformans* or viral infections, e.g. viruses that require host cell enzymes to synthesize and properly fold their viral envelope glycoproteins, or viruses that acquire a component of their envelope from an internal host cell membrane. GCS inhibition may result in improper glycoprocessing or the misfolding of one or more viral envelope glycoproteins, inhibition of viral secretion, or improper viral fusion of the virus with its target cells. Suitable viral infections for treatment
45 may be caused by, for example but not limited to, the following viruses: flaviviruses and pestiviruses, e.g. hepatitis C virus, yellow fever virus, dengue viruses 1-4, Japanese encephalitis virus, Murray Valley

encephalitis virus, Rocio virus, West Nile fever virus, St. Louis encephalitis virus, tick-borne encephalitis virus, Louping ill virus, Powassan virus, Omsk hemorrhagic fever virus, and Kyasanur forest disease virus; hepadnavirus, e.g. hepatitis B virus; paramyxovirus, e.g. respiratory syncytial virus or retroviruses, such as, human immunodeficiency virus.

5 The compounds of the present invention can also be used in the treatment of diseases in which excessive glycolipid synthesis occurs, such as, but not limited to, atherosclerosis, polycystic kidney disease and diabetic renal hypertrophy.

The compounds of the present invention can also be used in the treatment of neuronal disorders, such as, Alzheimer's disease or epilepsy; and neuronal degenerative diseases, such as, Parkinson's disease.

10 The compounds of the present invention can also be used in the treatment of neuronal injury, such as, spinal cord injuries or stroke.

The compounds of the present invention can also be used for reversibly rendering a male mammal infertile.

The compounds of the present invention can also be used in the treatment of obesity.

15 The compounds of the present invention can also be used in the treatment of inflammatory diseases or disorders associated with macrophage recruitment and activation, including but not limited to, rheumatoid arthritis, Crohn's disease, asthma and sepsis.

In additional aspects, therefore, the present invention provides:

20 (i) the use of a compound of formula (I) in the manufacture of a medicament for use as an inhibitor of glucosylceramide synthase.

(ii) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of a glycolipid storage disease. Examples of glycolipid storage disease which can be treated include, but are not limited to, Gaucher disease, Sandhoffs disease, Tay-Sachs disease, Fabry disease or GM1 gangliosidosis.

25 (iii) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of Niemann-Pick disease, types A and C.

(iv) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of mucopolysaccharidosis type I, mucopolysaccharidosis type IIIA, mucopolysaccharidosis type IIIB, mucopolysaccharidosis type VI or mucopolysaccharidosis type VII. Preferably the compounds are used in the
30 treatment of mucopolysaccharidosis type I.

(v) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of α -mannosidosis or mucopolipidosis type IV.

(vi) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of cancer in which glycolipid synthesis is abnormal, including but not limited to brain cancer, neuronal cancer,
35 neuroblastoma, renal adenocarcinoma, malignant melanoma, multiple myeloma and multi-drug resistant cancers.

(vii) the use of a compound of formula (I) in the manufacture of a medicament for use in the treatment of Alzheimer's disease, epilepsy or stroke.

(viii) the use of a compound of formula (I) in the manufacture of a medicament for use in the
40 treatment of Parkinson's disease.

(ix) the use of the compound of formula (I) in the manufacture of a medicament in the treatment of spinal injury.

(x) the use of a compound of formula (I) in the manufacture of a medicament for use in the treatment of diseases caused by infectious microorganisms which utilize glycolipids on the surface of cells as receptors
45 for either the organism itself or for toxins produced by the organism.

(xi) the use of a compound of formula (I) in the manufacture of a medicament for use in the treatment of disease caused by infectious organisms for which the synthesis of glucosylceramide is an essential or important process, such as but not limited to, pathologies associated with infections of pathogenic fungi, e.g. *Cryptococcus neoformans* or pathologies associated with viral infections.

5 (xii) the use of a compound of formula (I) in the manufacture of a medicament for use in the treatment of diseases associated with abnormal glycolipid synthesis, including but not limited to, polycystic kidney disease, diabetic renal hypertrophy and atherosclerosis.

(xiii) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of a condition treatable by the administration of a ganglioside, such as GM1 ganglioside. Examples of
10 such conditions are Parkinson's disease, stroke and spinal cord injuries.

(xiv) the use of a compound of formula (I) in the manufacture of a medicament for reversibly rendering a male mammal infertile.

(xv) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of obesity, e.g. as an appetite suppressant.

15 (xvi) the use of a compound of formula (I) in the manufacture of a medicament for the treatment of inflammatory diseases or disorders associated with macrophage recruitment and activation, including but not limited to, rheumatoid arthritis, Crohn's disease, asthma and sepsis.

(xvii) a method for the treatment of a glycolipid storage disease, e.g. Gaucher's disease, Sandhoffs disease, Tay-Sachs disease or GM1 gangliosidosis, which comprises the step of administering to a patient an
20 effective amount of a compound of formula (I).

(xviii) a method for the treatment of Niemann-Pick disease, types A and C, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xix) a method for the treatment of mucopolysaccharidosis type I, mucopolysaccharidosis type IIIA, mucopolysaccharidosis type IIIB, mucopolysaccharidosis type VI or mucopolysaccharidosis type
25 VII which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xx) a method for the treatment of α -mannosidosis or mucopolipidosis type IV which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxi) a method for the treatment of cancer in which glycolipid synthesis is abnormal, including but not limited to brain cancer, neuronal cancer, renal adenocarcinoma, malignant melanoma, multiple
30 myeloma and multi-drug resistant cancers, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxii) a method for the treatment of Alzheimer's disease, epilepsy or stroke which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxiii) a method for the treatment of Parkinson's disease, which comprises the step of administering
35 to a patient an effective amount of a compound of formula (I).

(xxiv) a method for the treatment of spinal injury which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxv) a method for the treatment of diseases caused by infectious microorganisms, which utilize glycolipids on the surface of cells as receptors for either the organism itself or for toxins produced by the
40 organism, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxvi) a method for the treatment of diseases caused by infectious organisms, e.g. pathogenic fungi or viruses, for which the synthesis of glucosylceramide is an essential or important process, such as but not limited to, pathologies associated with *Cryptococcus neoformans* infection, or pathologies associated with
45 viral infections, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxvii) a method for the treatment of diseases associated with abnormal glycolipid synthesis including but not limited to polycystic kidney disease, diabetic renal hypertrophy and atherosclerosis, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxviii) a method for the treatment of a condition treatable by the administration of a ganglioside, such as GM1 ganglioside, which comprises the step of administering to a patient an effective amount of a compound of formula (I). Examples of such conditions are, Parkinson's disease, stroke and spinal cord injuries.

(xxix) a method for reversibly rendering a male mammal infertile, which comprises the step of administering to said male mammal an effective amount of a compound of formula (I).

(xxx) a method for the treatment of obesity, which comprises the step of administering to a patient an effective amount of a compound of formula (I).

(xxxi) a method for the treatment of inflammatory diseases or disorders associated with macrophage recruitment and activation, including but not limited to, rheumatoid arthritis, Crohn's disease, asthma and sepsis, which comprises the step of administering to a patient an effective amount of a compound of formula (I).


The invention also provides for the use of a compound of formula (I) for the treatment of the above mentioned diseases and conditions.

All publications, including, but not limited to, patents and patent applications, cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The invention will now be described by reference to the following examples, which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.


Examples

Example 1 Benzamide, *N*-[(4-fluorophenyl)methyl]-4-[[[(2*S*,3*S*,4*R*,5*S*)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2*S*,3*S*,4*R*,5*S*) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added *N*-[(4-fluorophenyl)methyl]-4-formylbenzamide (196mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g), which had been pre-washed with 10% aqueous hydrochloric acid. The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid (70mg, 57%). ¹H NMR (d₄-methanol) δ 2.52 (1H, m), 2.65 (1H, m), 2.78 (1H, m), 3.60-4.16 (7H, m), 4.5-4.7 (2H, m), 7.06 (2H, m), 7.38 (2H, m), 7.52 (2H, d, *J* = 8 Hz), 7.82 (2H, d, *J* = 8 Hz). MS *m/z* 404.9 (M)⁺.

Example 2 3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[[4-(phenylmethoxy)phenyl]methyl]-, (2S,3S,4R,5S)




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)- (2S,3S,4R,5S) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added (4-phenylmethoxy)benzaldehyde (162mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g), which had been pre-washed with 10% aqueous hydrochloric acid. The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid (30mg, 27%). ¹H NMR (d4-methanol) δ 2.52 (1H, m), 2.64 (1H, m), 2.75 (1H, m), 3.53-4.0 (7H, m), 5.15 (2H, s), 6.95 (2H, d, J = 8.3 Hz), 7.25-7.48 (7H, m). MS *m/z* 360.0 (M+H)⁺.

Example 3 Benzamide, N-[1-(S)-(phenyl)ethyl]-4-[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2S,3S,4R,5S) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added *N*-[1-(S)-(phenyl)ethyl]-4-formylbenzamide (193mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g), which had been pre-washed with 10% aqueous hydrochloric acid. The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid (10mg, 8%). ¹H NMR (d4-methanol) δ 1.51 (3H, d, J = 7.1 Hz), 2.53 (1H, dd, J = 6.8 and 12.1 Hz), 2.66 (1H, dd, J = 3.8 and 12.1 Hz), 2.79 (1H, m), 3.62-4.00 (6H, m), 4.13 (1H, d, J = 13.6 Hz), 5.25 (1H, m), 7.24 (1H, t, J = 7.2 Hz), 7.3-7.4 (4H, m), 7.52 (2H, d, J = 8.3 Hz), 7.81 (2H, d, J = 8.3 Hz). MS *m/z* 401.0 (M+1)⁺.

Example 4 3,4,5-Piperidinetriol, 1-[(3-cyano-4-(dipropylamino)phenyl)methyl]-2-(hydroxymethyl)-, (2S,3S,4R,5S)




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2S,3S,4R,5S) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added 3-cyano-4-(dipropylamino)benzaldehyde (162mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g), which had been pre-washed with 10% aqueous hydrochloric acid. The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was then eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid (30mg, 27%). ¹H NMR (d₄-methanol) δ 0.93 (6H, t, J = 7.3 Hz), 1.6 (4H, m), 2.5-2.85 (3H, m), 3.34 (4H, m), 3.55-4.05 (7H, m), 7.03 (1H, d, J = 9 Hz), 7.5 (1H, dd, J = 1.9 and 9 Hz), 7.61 (1H, d, J = 1.9 Hz). MS *m/z* 378.1 (M+H)⁺.

Example 5 Benzamide, N-[1-(S)-(4-fluorophenyl)ethyl]-4-[[[(2S,3S,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidiny]methyl]-,




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2S,3S,4R,5S) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added *N*-[1-(S)-(4-fluorophenyl)ethyl]-4-formylbenzamide (207mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g), which had been pre-washed with 10% aqueous hydrochloric acid. The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was then eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid, 30mg (24%). ¹H NMR (d₄-methanol) δ 1.57 (3H, d, J = 6.8 Hz), 2.54 (1H, dd, J = 6.4 and 12.1 Hz), 2.65 (1H, dd, J = 3.8 and 12.1 Hz), 2.79 (1H, m), 3.6-3.98 (6H, m), 4.12 (1H, d, J = 14 Hz), 5.23 (1H, q, J = 7.2 Hz), 7.06 (2H, m), 7.43 (2H, m), 7.52 (2H, d, J = 8.3 Hz), 7.81 (2H, d, J = 8.3 Hz). MS *m/z* 419.0 (M+H)⁺.

Example 6 Benzamide, *N*-[1-(*R*)-(phenyl)ethyl]-4-[[[(2*S*,3*S*,4*R*,5*S*)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2*S*,3*S*,4*R*,5*S*) (150mg, 0.92mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (540mg, 2.3mmol) in methanol (5ml) was added *N*-[1-(*R*)-(phenyl)ethyl]-4-formylbenzamide (910mg, 3.59mmol). Dichloromethane (1ml) was added to dissolve the aldehyde and the mixture warmed to dissolve the amine. Acetic acid (0.1ml, 1.75mmol) was added to the reaction mixture which was then stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (5g) (which had been pre-washed with 10% aqueous hydrochloric acid, water and then methanol). The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was then eluted using a solution of 4:1 methanol/ammonium hydroxide (50ml), then 3:1:1 methanol/dichloromethane/ammonium hydroxide (50ml). The resulting solution was concentrated to give a gum which was dissolved in hot water, cooled and freeze dried to afford the title compound as a white solid (300mg 81%). ¹H NMR (d4-methanol) δ 1.56 (3H, d, *J* = 7.0 Hz), 2.51 (1H, dd, *J* = 6.8 and 12.0 Hz), 2.64 (1H, dd, *J* = 3.8 and 12.0 Hz), 2.77 (1H, ddd, *J* = 4.5, 5.3 and 6.0 Hz), 3.64 (1H, dd, *J* = 3.4 and 6.4 Hz), 3.67 (1H, d, *J* = 13.9 Hz), 3.73 (1H, ddd, *J* = 3.8, 6.4 and 6.8 Hz), 3.83 (1H, dd, *J* = 5.3 and 11.7 Hz), 3.91 (1H, dd, *J* = 4.5 and 11.7 Hz), 3.96 (1H, dd, *J* = 3.4 and 6.0 Hz), 4.12 (1H, d, *J* = 13.9 Hz), 5.24 (1H, q, *J* = 7.0 Hz), 7.22 (1H, t, *J* = 7.2 Hz), 7.32 (2H, t, *J* = 7.2 Hz), 7.39 (2H, d, *J* = 7.2 Hz), 7.50 (2H, d, *J* = 8.3 Hz), 7.80 (2H, d, *J* = 8.3 Hz). MS *m/z* 401.2 (M+H)⁺.

Example 7 Benzamide, *N*-[1-(*R*)-(4-fluorophenyl)ethyl]-4-[[[(2*S*,3*S*,4*R*,5*S*)-3,4,5-trihydroxy-2-(hydroxymethyl)-1-piperidinyl]methyl]-,




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2*S*,3*S*,4*R*,5*S*) (50mg, 0.31mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (178mg, 0.76mmol) in 10% acetic acid in methanol (2ml) was added *N*-[1-(*R*)-(4-fluorophenyl)ethyl]-4-formylbenzamide (207mg, 0.76mmol) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (3g) (which had been pre-washed with 10% aqueous hydrochloric acid). The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was then eluted using a solution of 2:2:1 methanol/water/ammonium hydroxide (100ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried to afford the title compound as a white solid, 40mg (31%). ¹H NMR (d4-methanol) δ 1.55 (3H, d, *J* = 7.1 Hz), 2.52 (1H, dd, *J* = 6.8 and 12.0 Hz), 2.64 (1H, dd, *J* = 3.8 and 12.0 Hz), 2.77 (1H, ddd, *J* = 4.5, 5.1 and 5.9 Hz), 3.64 (1H, dd, *J* = 3.4 and 6.4 Hz), 3.67 (1H, d, *J* = 14 Hz), 3.74 (1H, ddd, *J* = 3.8, 6.4 and 6.8 Hz), 3.83 (1H, dd, *J* = 5.1 and 11.5 Hz), 3.91 (1H, dd, *J* = 4.5 and 11.5 Hz), 3.96 (1H, dd, *J* = 3.4 and 6.0 Hz), 4.12 (1H, d, *J* = 14 Hz), 5.22 (1H, q, *J* = 7.1 Hz), 7.05 (2H, dd, *J* = 8.7 and 8.9 Hz), 7.41 (2H, dd, *J* = 5.6 and 8.7 Hz), 7.50 (2H, d, *J* = 8.1 Hz), 7.79 (2H, d, *J* = 8.1 Hz). MS *m/z* 419.1 (M+H)⁺.

Example 8 3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[(2-phenyl-2H-1,4-benzoxazin-3(4H)-one-6-yl)methyl]-, (2S,3S,4R,5S)




To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)-, (2S,3S,4R,5S) (50mg, 0.31mmol), 2-phenyl-2H-1,4-benzoxazin-3(4H)-one-6-carbaldehyde (230mg, 0.91mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (300mg, 1.31mmol) in methanol (2ml) was added acetic acid (0.2ml). The resultant mixture was stirred at room temperature for 16h. The crude reaction mixture was purified using a plug of acidic Dowex 50X4-200 resin (1g) (which had been pre-washed with methanol (10ml)). The resin was eluted with methanol (25ml) to remove all non-basic side products. The desired compound was then eluted using a solution of 7:1 methanol/ammonium hydroxide (25ml). The resulting solution was concentrated to a small volume (1ml) and freeze dried. The freeze dried solid was further purified using silica chromatography (gradient elution 10 to 30% methanol/dichloromethane and 1% ammonia) to afford the title compound as a white solid (34mg, 27%).
¹H NMR (d4-methanol) δ 2.62 (1H, m), 2.75 (1H, m), 2.87 (1H, m), 3.66-4.11 (7H, m), 5.69 (1H, s), 6.94-7.05 (3H, m), 7.32-7.45 (5H, m). MS *m/z* 401.2 (M+H)⁺.

Example 9 3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[[4-[(4-chlorophenyl)methoxy]phenyl]methyl]-, (2S,3S,4R,5S)



To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)- (2S,3S,4R,5S) (57mg, 0.35mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (253mg, 1.08mmol) in 10% acetic acid in methanol (2ml) was added 4-[(4-chlorophenyl)methoxy] benzaldehyde (280mg, 1.13mmol) and the reaction mixture was stirred at room temperature for two days. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (1g), which had been pre-washed with methanol. The resin was eluted with methanol (25ml) and acetone (100ml) to remove all non-basic side products. The desired compound was eluted using a solution of 7:1 methanol/ammonium hydroxide (50ml). The resulting solution was concentrated, re-dissolved in methanol and pre-adsorbed onto silica gel. The product was then eluted through a silica gel column using methanol as solvent, concentrated to a gum, and freeze dried to afford the title compound as a white solid (64mg, 46%).
¹H NMR (d4-methanol) δ 2.51 (1H, m), 2.63 (1H, m), 2.74 (1H, m), 3.53-4.0 (7H, m), 5.05 (2H, s), 6.93 (2H, d, J = 8.7 Hz), 7.29 (2H, d, J = 8.7 Hz), 7.37 (2H, d, J = 8.5 Hz), 7.43 (2H, d, J = 8.5 Hz). MS *m/z* 394.1, 396.1 (M+H)⁺.

Example 10 3,4,5-Piperidinetriol, 2-(hydroxymethyl)-1-[[4-[(4-fluorophenyl)methoxy]phenyl]methyl]-, (2S,3S,4R,5S)



To a mixture of 3,4,5-piperidinetriol, 2-(hydroxymethyl)- (2S,3S,4R,5S) (57mg, 0.35mmol) and (polystyrylmethyl)trimethylammonium cyanoborohydride (250mg, 1.07mmol) in 10% acetic acid in methanol (2ml) was added 4-[(4-fluorophenyl)methoxy] benzaldehyde (250mg, 1.08mmol) and the reaction mixture was stirred at room temperature for two days. The reaction mixture was purified using a plug of acidic Dowex 50WX-12 resin (1g), which had been pre-washed with methanol. The resin was eluted with methanol (25ml) and ethyl acetate (100ml) to remove all non-basic side products. The desired compound was eluted using a solution of 7:1 methanol/ammonium hydroxide (100ml). The resulting solution was concentrated, re-dissolved in methanol and pre-adsorbed onto silica gel. The product was then eluted through a silica gel column using methanol as solvent, concentrated to a gum, and freeze dried to afford the title compound as a white solid (56mg, 42%). ¹H NMR (d₄-methanol) δ 2.57 (1H, m), 2.69 (1H, m), 2.80 (1H, m), 3.60-4.10 (7H, m), 5.05 (2H, s), 6.95 (2H, d, J = 8.5 Hz), 7.09 (2H, m), 7.31 (2H, d, J = 8.5 Hz), 7.45 (2H, m). MS *m/z* 378.1 (M+H)⁺.

Biological Assays

The compounds of the invention may be tested for their biological activity in the following assays:

Inhibition of GCS

The assay for inhibition of GCS was performed essentially as described in Platt et al., J.Biol.Chem., (1994), 269, 27108, the enzyme source being human recombinant GCS expressed in insect cells.

Inhibition of non-lysosomal-β-glucocerebrosidase

The assay for inhibition of non-lysosomal-β-glucocerebrosidase was essentially carried out as described in Overkleeft, H. S. et al., J. Biol. Chem., (1998) 273, 26522-26527 with the following differences: whole cell extracts of MCF7 (a human breast carcinoma cell line) was used as the source of the enzyme instead of splenic membrane suspensions; 5mM instead of 3Mm, 4-MU β-glucoside was used as substrate and 0.2M citrate/phosphate (pH 5.8) was used instead of McIlvaine buffer.

Table I shows IC₅₀ data for compounds of the invention against human GCS and non-lysosomal-β-glucocerebrosidase enzymes.

Table I

Compound	Inhibition of GCS (IC ₅₀ μM)	Inhibition of non-lysosomal-β-glucocerebrosidase (IC ₅₀ μM)
Example 1	2.20	100.00
Example 2	0.66	9.40
Example 3	7.20	14.90
Example 4	0.62	1.93
Example 7	4.70	>100.00

Estimating the cell-based IC_{50} for GCS inhibition by measuring glucosylceramide (GlcCer) depletion

Human mammary epithelial cells (MCF-7) were cultured for 5-7 days, with varying concentrations of a compound of the invention to be tested (0; 0.01; 0.05; 0.25; 1.25 and 6.25 μ M). The cells were harvested and the total cellular lipids extracted. Neutral glycolipids were separated by partitioning in a DIPE/1-
5 butanol/saline suspension, according to methods well known to those skilled in the art. The neutral glycolipid extracts were then separated by High-Performance Thin Layer Chromatography (HPTLC), using non-polar TLC conditions (chloroform: methanol: 0.2% $CaCl_2$; 65:35:8), according to methods well known to those skilled in the art. GlcCer bands were visualized and the TLC plates were scanned immediately. Scion Image software was then used to quantify GlcCer in the samples relative to a GlcCer standard. This enabled a cell-
10 based IC_{50} to be calculated for compounds of the invention for GCS inhibition, as shown in the Table II.


Table II shows cell-based IC_{50} data for compounds of the invention against human GCS.

Table II

Compound	Inhibition of GCS Cellular-based (IC_{50} μ M)
Example 2	< 0.10
Example 6	1.00

CLAIMS:

1. A compound of formula (I) or a pharmaceutically acceptable salt or prodrug thereof:



wherein

R is C_{1-3} alkylAr¹ where Ar¹ is phenyl or pyridyl;

wherein phenyl is substituted by one or more substituents selected from CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R⁵)₂, N(R¹)COR², N(R¹)SO_nR², C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one or more of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided, that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ phenyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O, S and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C₁₋₆ alkyl and C₀₋₃ alkylAr⁴;

and the Ar¹ phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF₃, OCF₃, OR³ and C₁₋₆ alkyl;

and wherein pyridyl is substituted by one or more substituents, selected from, CN, CON(R¹)₂, SO₂R², SO₂N(R¹)₂, N(R⁵)₂, N(R¹)COR², N(R¹)SO_nR², F, Cl, Br, CF₃, OCF₃, OR³, C₁₋₆ alkyl, C₀₋₆ alkylAr², C₂₋₆ alkenylAr² and C₃₋₆ alkynylAr² wherein one of the -CH₂- groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR³, provided that when the heteroatom is O, at least two -CH₂- groups separate it from any additional O atom in the alkyl chain; or two adjacent substituents on the Ar¹ pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O, S and NR⁴ and is optionally substituted by one or more substituents selected from, an oxo group, C₁₋₆ alkyl and C₀₋₃ alkylAr⁴;

R¹ is H, C₁₋₆ alkyl optionally substituted by OH, Ar³, or C₁₋₆ alkylAr³, or the group N(R¹)₂ may form a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O, S and NR³ and is optionally substituted by an oxo group;

R² is C₁₋₆ alkyl optionally substituted by OH, Ar³, or C₁₋₆ alkylAr³;

R³ is H, or C₁₋₆ alkyl;

R⁴ is H, C₁₋₆ alkyl or C₀₋₃ alkylAr⁴;

R⁵ is H, C₁₋₆ alkyl optionally substituted by OH, Ar³, or C₁₋₆ alkylAr³, or the group N(R⁵)₂ may form a 5- to 10-membered heterocyclic group optionally containing one or more additional heteroatoms selected from O, S and NR³ and is optionally substituted by an oxo group;


Ar² and Ar³ are independently phenyl or a 5- to 10-membered heteroaryl group containing up to 3 heteroatoms selected from O, S and NR³, which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF₃, OCF₃, OR³ and C₁₋₆ alkyl;

Ar⁴ is phenyl or pyridyl either of which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF₃, OCF₃, OR³ and C₁₋₆ alkyl; and

n = 0, 1 or 2.


2. A compound as defined in claim 1 wherein R is C₁ alkylAr¹.

3. A compound as defined in claim 1 or 2 wherein Ar^1 is phenyl, wherein phenyl is substituted as defined for claim 1.
4. A compound as defined in any one of the preceding claims wherein Ar^1 is phenyl, wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, $N(R^5)_2$, and $C_{0-6}alkylAr^2$ wherein one or more of the $-CH_2-$ groups of the alkyl chain may be replaced with a heteroatom selected from O, S and NR^3 , provided that when the heteroatom is O, at least two $-CH_2-$ groups separate it from any additional O atom in the alkyl chain, or two adjacent substituents on the Ar^1 pyridyl may together form a fused 5- or 6-membered saturated or unsaturated ring wherein the ring optionally contains 1 or 2 heteroatoms selected from O and NR^4 and is optionally substituted by one or more substituents selected from, an oxo group, $C_{1-6}alkyl$ and $C_{0-3}alkylAr^4$, and the Ar^1 phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , OR^3 and $C_{1-6}alkyl$.
5. A compound as defined in any one of the preceding claims wherein Ar^1 is phenyl, wherein phenyl is substituted by one or more substituents selected from CN, $CON(R^1)_2$, $N(R^5)_2$, and $C_{0-6}alkylAr^2$ wherein one or more of the $-CH_2-$ groups of the alkyl chain may be replaced with O, provided that at least two $-CH_2-$ groups separate it from any additional O atom introduced into the alkyl chain and the Ar^1 phenyl is optionally substituted by one or more additional substituents selected from F, Cl, Br, CF_3 , OCF_3 , OR^3 and $C_{1-6}alkyl$.
6. A compound as defined in any one of the preceding claims wherein Ar^2 is phenyl which is optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and $C_{1-6}alkyl$.
7. A compound as defined in any one of the preceding claims wherein R^1 is H or $C_{1-6}alkylAr^3$.
8. A compound as defined in any one of the preceding claims wherein R^4 is H or $C_{1-6}alkyl$.
9. A compound as defined in any one of the preceding claims wherein Ar^3 is phenyl which may be optionally substituted by one or more substituents selected from F, Cl, Br, CN, CF_3 , OCF_3 , OR^3 and $C_{1-6}alkyl$.
10. A compound as defined in any one of the preceding claims wherein R^5 is $C_{1-6}alkyl$.
11. A compound of formula (I) as described in any one of Examples 1 to 10 or a pharmaceutically acceptable salt or prodrug thereof.
12. A compound as defined in any one of the preceding claims for use in medicine.
13. A pharmaceutical composition comprising a compound of formula (I) as defined in any one of claims 1 to 11, together with one or more pharmaceutically acceptable carriers, excipients and/or diluents.
14. A process for the preparation of a compound of formula (I) as defined in any one of claims 1 to 11 which comprises:
- a) reductive amination of an aldehyde of formula R^5CHO wherein R^5 is $C_{0-2}alkylAr^1$ where Ar^1 is as defined in claim 1, with a compound of formula (II):



or

b) deprotection of a compound of formula (III):



wherein R is as defined in claim 1 and P, which may be the same or different, are hydroxy protecting groups.

- 10 15. The use of a compound of formula (I) as defined in any one of claims 1 to 11 in the manufacture of a medicament for use as an inhibitor of glucosylceramide synthase.
16. The use of a compound as defined in any one of claims 1 to 11 in the manufacture of a medicament for the treatment of a glycolipid storage disease.
- 15 17. The use as claimed in claim 16 wherein the glycolipid storage disease is Gaucher disease, Sandhoffs disease, Tay-Sachs disease, Fabry disease or GM1 gangliosidosis.
18. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for the treatment of Niemann-Pick disease type C, mucopolysaccharidosis type I, mucopolysaccharidosis type IIIA, mucopolysaccharidosis type IIIB, mucopolysaccharidosis type VI, mucopolysaccharidosis type VII, α -mannosidosis or mucolipidosis type IV.
- 20 19. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for the treatment of cancer in which glycolipid synthesis is abnormal.
- 25 20. The use as claimed in claim 19 wherein the cancer in which glycolipid synthesis is abnormal is selected from, brain cancer, neuronal cancer, neuroblastoma, renal adenocarcinoma, malignant melanoma, multiple myeloma and multi-drug resistant cancer.
- 30 21. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for use in the treatment of Alzheimer's disease, epilepsy, stroke, Parkinson's disease or spinal injury.
22. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for use in the treatment of diseases caused by, infectious microorganisms which utilize glycolipids on the surface of cells as receptors for either the organism itself or for toxins produced by the organism, or infectious organisms for which the synthesis of glucosylceramide is an essential or important process.
- 35 23. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for use in the treatment of diseases associated with abnormal glycolipid synthesis.
- 40

24. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for the treatment of a condition treatable by the administration of a ganglioside.

5 25. The use as claimed in claim 24, wherein the condition is treatable by the administration of a GM1 ganglioside.

26. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for use in reversibly rendering a male mammal infertile.

10

27. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for the treatment of obesity.


15

28. The use of a compound as defined in any one of claims 1 to 11, in the manufacture of a medicament for the treatment of inflammatory diseases or disorders associated with macrophage recruitment and activation.

29. The use as claimed in claim 28, wherein the inflammatory disease or disorder associated with macrophage recruitment and activation is selected from rheumatoid arthritis, Crohn's disease, asthma and sepsis.

20

30. A compound of formula (III):



25 wherein R is as defined in claim 1 and P, which may be the same or different, are hydroxy protecting groups.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 03/03244

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D211/46 C07D405/06 C07D401/06 C07D413/06 A61K31/445 A61P3/00 A61P35/00 A61P9/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data, INSPEC		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	WO 02 055498 A (BUTTERS TERENCE D ;OXFORD GLYCOSCIENCES UK LTD (GB); UNIV OXFORD () 18 July 2002 (2002-07-18) cited in the application claims page 21 -page 22; examples 9,10 page 24 -page 26; examples 13-15 --- -/--	1-30
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : 'A' document defining the general state of the art which is not considered to be of particular relevance 'E' earlier document but published on or after the international filing date 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) 'O' document referring to an oral disclosure, use, exhibition or other means 'P' document published prior to the international filing date but later than the priority date claimed 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art 'G' document member of the same patent family		
Date of the actual completion of the international search 11 November 2003		Date of mailing of the international search report 20/11/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer Stix-Malaun, E

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/GB 03/03244

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BARILI, PIER LUIGI ET AL: "Rare and complex saccharides from D-galactose and other milk derived carbohydrates. 7. Double reductive amination of L-arabino-hexos-5- uloses: a diastereoselective approach to 1-deoxy-D-galactostatin derivatives" TETRAHEDRON (1997), 53(9), 3407-3416 , XP004105422 abstract page 3409; examples 14-17 page 3407, paragraph 1</p>	1-30
A	<p>MEHTA G ET AL: "A norbornyl route to azasugars: a new synthesis of deoxynojirimycin analogues" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 41, no. 30, 22 July 2000 (2000-07-22), pages 5741-5745, XP004209550 ISSN: 0040-4039 abstract page 5743; example 17</p>	1-30
A	<p>MERRER LE Y ET AL: "SYNTHESIS OF AZASUGARS AS POTENT INHIBITORS OF GLYCOSIDASES" BIOORGANIC & MEDICINAL CHEMISTRY, ELSEVIER SCIENCE LTD, GB, vol. 5, no. 3, March 1997 (1997-03), pages 519-533, XP001062185 ISSN: 0968-0896 abstract page 521; table I</p>	1-30
A	<p>POITOUT L ET AL: "Synthesis of Azasugars. Part 1 Isomerization of Polyhydroxylated Piperidines" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 37, no. 10, 4 March 1996 (1996-03-04), pages 1609-1612, XP004030006 ISSN: 0040-4039 abstract page 1609; examples 2A-4A</p>	1-30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 03/03244

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-30(part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 03 03244

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-30(part)

Present claims 1-30 relate to compounds defined by reference to a desirable characteristic or property, namely "prodrug" and "protecting group".

The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compounds by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds according to formulae (I) or (III). With respect to the compounds of formula (III) the structural definitions of the description have been taken into account.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internati Application No

PCT/GB 03/03244

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 02055498 A	18-07-2002	CA 2433675 A1	18-07-2002
		WO 02055498 A1	18-07-2002
		WO 02055064 A2	18-07-2002